


Atoms, elements and compounds

Atom	<i>The smallest part of an element that can exist</i>	Have a radius of around 0.1 nanometres and have no charge (0).
Element	<i>Contains only one type of atom</i>	Around 100 different elements each one is represented by a symbol e.g. O, Na, Br.
Compound	<i>Two or more elements chemically combined</i>	Compounds can only be separated into elements by chemical reactions.



Central nucleus	Contains protons and neutrons
Electron shells	Contains electrons

Name of Particle	Relative Charge	Relative Mass
Proton	+1	1
Neutron	0	1
Electron	-1	Very small

Electronic shell	Max number of electrons
1	2
2	8
3	8
4	2

Electronic structures

AQA GCSE Atomic structure and periodic table part 1

Relative electrical charges of subatomic particles

7 Li	Mass number	<i>The sum of the protons and neutrons in the nucleus</i>
3	Atomic number	<i>The number of protons in the atom</i> Number of electrons = number of protons

Mixtures	<i>Two or more elements or compounds not chemically combined together</i>	Can be separated by physical processes.
-----------------	---	---

Pre 1900		<i>Tiny solid spheres that could not be divided</i>	Before the discovery of the electron, John Dalton said the solid sphere made up the different elements.
1897 'plum pudding'		<i>A ball of positive charge with negative electrons embedded in it</i>	JJ Thompson 's experiments showed that showed that an atom must contain small negative charges (discovery of electrons).
1909 nuclear model		<i>Positively charge nucleus at the centre surrounded negative electrons</i>	Ernest Rutherford's alpha particle scattering experiment showed that the mass was concentrated at the centre of the atom.
1913 Bohr model		<i>Electrons orbit the nucleus at distances</i>	Niels Bohr proposed that electrons orbited in fixed shells; this was supported by experimental observations.

James Chadwick
Provided the evidence to show the existence of neutrons within the nucleus

Rutherford's scattering experiment

A beam of alpha particles are directed at a very thin gold foil

Most of the alpha particles passed right through. A few (+) alpha particles were deflected by the positive nucleus. A tiny number of particles reflected back from the nucleus.

Chemical equations	<i>Show chemical reactions - need reactant(s) and product(s) energy always involves and energy change</i>	Law of conservation of mass states the total mass of products = the total mass of reactants.
---------------------------	---	--

Word equations	<i>Uses words to show reaction</i> reactants → products magnesium + oxygen → magnesium oxide	Does not show what is happening to the atoms or the number of atoms.
Symbol equations	<i>Uses symbols to show reaction</i> reactants → products $2\text{Mg} + \text{O}_2 \rightarrow 2\text{MgO}$	Shows the number of atoms and molecules in the reaction, these need to be balanced.

Relative atomic mass	Isotopes	<i>Atoms of the same element with the same number of protons and different numbers of neutrons</i>	^{35}Cl (75%) and ^{37}Cl (25%) Relative abundance = (% isotope 1 x mass isotope 1) + (% isotope 2 x mass isotope 2) ÷ 100 e.g. (25 x 37) + (75 x 35) ÷ 100 = 35.5
-----------------------------	-----------------	--	---

Method	Description	Example
Filtration	<i>Separating an insoluble solid from a liquid</i>	To get sand from a mixture of sand, salt and water.
Crystallisation	<i>To separate a solid from a solution</i>	To obtain pure crystals of sodium chloride from salt water.
Simple distillation	<i>To separate a solvent from a solution</i>	To get pure water from salt water.
Fractional distillation	<i>Separating a mixture of liquids each with different boiling points</i>	To separate the different compounds in crude oil.
Chromatography	<i>Separating substances that move at different rates through a medium</i>	To separate out the dyes in food colouring.

Alkali metals

Transition metals

Haloge

Noble gases

1	2	3	4 ^{ns}	5	6	7	0										
H							He										
Li	Be		B	C	N	O	F	Ne									
Na	Mg		Al	Si	P	S	Cl	Ar									
K	Ca	Sc	Ti	V	Cr	Mn	Fe	Co	Ni	Cu	Zn	Ga	Ge	As	Se	Br	Kr
Rb	Sr	Y	Zr	Nb	Mo	Tc	Ru	Rh	Pd	Ag	Cd	In	Sn	Sb	Te	I	Xe
Cs	Ba	La	Hf	Ta	W	Re	Os	Ir	Pt	Au	Hg	Tl	Pb	Bi	Po	At	Rn
Fr	Ra	Ac	Rf	Db	Sg	Bh	Hs	Mt	?	?	?						

Elements arranged in order of atomic number

Elements with similar properties are in columns called groups

Elements in the same group have the same number of outer shell electrons and elements in the same period (row) have the same number of electron shells.

The Periodic table

Development of the Periodic table

Before discovery of protons, neutrons and electrons

Elements arranged in order of atomic weight

Early periodic tables were incomplete, some elements were placed in inappropriate groups if the strict order atomic weights was followed.

Mendeleev

Left gaps for elements that hadn't been discovered yet

Elements with properties predicted by Mendeleev were discovered and filled in the gaps. Knowledge of isotopes explained why order based on atomic weights was not always correct.

Metals	To the left of the Periodic table	Form positive ions. Conductors, high melting and boiling points, ductile, malleable.
Non metals	To the right of the Periodic table	Form negative ions. Insulators, low melting and boiling points.

Metals to the left of this line, non metals to the right

Metals and non metals

AQA GCSE Atomic structure and periodic table

Group 1

Alkali metals

Alkali metals	Very reactive with oxygen, water and chlorine	Only have one electron in their outer shell. Form +1 ions.
	Reactivity increases down the group	Negative outer electron is further away from the positive nucleus so is more easily lost.

Halogens	Consist of molecules made of a pair of atoms	Have seven electrons in their outer shell. Form -1 ions.
	Melting and boiling points increase down the group (gas → liquid → solid)	Increasing atomic mass number.
	Reactivity decreases down the group	Increasing proton number means an electron is more easily gained e.g. NaCl

Group

Transition metals (Chemistry only)

With oxygen	Forms a metal oxide	Metal + oxygen → metal oxide	e.g. $4\text{Na} + \text{O}_2 \rightarrow 2\text{Na}_2\text{O}$
With water	Forms a metal hydroxide and hydrogen	Metal + water → metal hydroxide + hydrogen	e.g. $2\text{Na} + 2\text{H}_2\text{O} \rightarrow 2\text{NaOH} + \text{H}_2$
With chlorine	Forms a metal chloride	Metal + chlorine → metal chloride	e.g. $2\text{Na} + \text{Cl}_2 \rightarrow 2\text{NaCl}$

Noble gases	Unreactive, do not form molecules	This is due to having full outer shells of electrons.
	Boiling points increase down the group	Increasing atomic number.


Compared to group 1	<ul style="list-style-type: none"> Less reactive Harder Denser Higher melting points 	<ul style="list-style-type: none"> Cu^{2+} is blue Ni^{2+} is pale green, used in the manufacture of margarine
Typical properties	<ul style="list-style-type: none"> Many have different ion possibilities with different charges Used as catalysts Form coloured compounds 	<ul style="list-style-type: none"> Fe^{2+} is green, used in the Haber process Fe^{3+} is reddish-brown Mn^{2+} is pale pink

With metals	Forms a metal halide	Metal + halogen → metal halide e.g. Sodium + chlorine → sodium chloride	metal atom loses outer shell electrons and halogen gains an outer shell electron
With hydrogen	Forms a hydrogen halide	Hydrogen + halogen → hydrogen halide e.g. Hydrogen + bromine → hydrogen bromide	e.g. $\text{Cl}_2 + \text{H}_2 \rightarrow 2\text{HCl}$
With aqueous solution of a halide salt	A more reactive halogen will displace the less reactive halogen from the salt	Chlorine + potassium bromide → potassium chloride + bromine	e.g. $\text{Cl}_2 + 2\text{KBr} \rightarrow 2\text{KCl} + \text{Br}_2$

Ionic	Particles are oppositely charged ions	Occurs in compounds formed from metals combined with non metals.
Covalent	Particles are atoms that share pairs of electrons	Occurs in most non metallic elements and in compounds of non metals.
Metallic	Particles are atoms which share delocalised electrons	Occurs in metallic elements and alloys.

Solid, liquid, gas

Melting and freezing happen at melting point, boiling and condensing happen at boiling point.



(HT only)
Limitations of simple model:

- There are no forces in the model
- All particles are shown as spheres
- Spheres are solid

s	solid
l	liquid
g	gas

Chemical bonds

The three states of matter

Good conductors of electricity

Delocalised electrons carry electrical charge through the metal.

Good conductors of thermal energy

Energy is transferred by the delocalised electrons.

High melting and boiling points

This is due to the strong metallic bonds.

Pure metals can be bent and shaped

Atoms are arranged in layers that can slide over each other.

AQA BONDING, STRUCTURE AND THE PROPERTIES OF MATTER 1

High melting and boiling points

Large amounts of energy needed to break the bonds.

Do not conduct electricity when solid

Ions are held in a fixed position in the lattice and cannot move.

Do conduct electricity when molten or dissolved

Lattice breaks apart and the ions are free to move.

Properties of ionic compounds

Metals as conductors

Properties of metals and alloys

Alloys

Mixture of two or more elements at least one of which is a metal

Harder than pure metals because atoms of different sizes disrupt the layers so they cannot slide over each other.

Ionic bonding

Electrons are transferred so that all atoms have a noble gas configuration (full outer shells).



Metal atoms lose electrons and become positively charged ions

Group 1 metals form +1 ions
Group 2 metals form +2 ions


Non metals atoms gain electrons to become negatively charged ions

Group 6 non metals form -2 ions
Group 7 non metals form -1 ions

Metallic bonding



Ionic compounds



Structure


- Held together by strong electrostatic forces of attraction between oppositely charged ions
- Forces act in all directions in the lattice

Giant structure of atoms arranged in a regular pattern

Delocalised electrons

Metal ions

Electrons in the outer shell of metal atoms are delocalised and free to move through the whole structure. This sharing of electrons leads to strong metallic bonds.

Very large molecules	<i>Solids at room temperature</i>	Atoms are linked by strong covalent bonds.	$\begin{array}{c} \text{H} & \text{H} \\ & \\ \text{C} = & \text{C} \\ & \\ \text{H} & \text{H} \end{array} \rightarrow \left(\begin{array}{c} \text{H} & \text{H} \\ & \\ \text{---C} & \text{---C} \\ & \\ \text{H} & \text{H} \end{array} \right)_n$	<i>Each carbon atom is bonded to four others</i>		Very hard.	Rigid structure.
						Very high melting point.	Strong covalent bonds.
						Does not conduct electricity.	No delocalised electrons.


Usually gases or liquids	<i>Covalent bonds in the molecule are strong but forces between molecules (intermolecular) are weak</i>	Low melting and boiling points.	Due to having weak intermolecular forces that easily broken.
		Do not conduct electricity.	Due to them molecules not having an overall electrical charge.
		Larger molecules have higher melting and boiling points.	Intermolecular forces increase with the size of the molecules.

Properties of small molecules	Polymers	Diamond
	AQA BONDING, STRUCTURE AND THE PROPERTIES OF MATTER 2	
	Size of particles and their properties (Chemistry only)	
	Nanoparticles <i>Between 1 and 100 nanometres (nm) in size</i>	1 nanometre (1 nm) = 1×10^{-9} metres (0.000 000 001m or a billionth of a metre).

Diamond, graphite, silicon dioxide	<i>Very high melting points</i>	Lots of energy needed to break strong, covalent bonds.
------------------------------------	---------------------------------	--


Atoms share pairs of electrons

Can be small molecules e.g. ammonia




Dot and cross :
+ Show which atom the electrons in the bonds come from
- All electrons are identical


2D with bonds:
+ Show which atoms are bonded together
- It shows the H-C-H bond incorrectly at 90°




Can be giant covalent structures e.g. polymers





3D ball and stick model:
+ Attempts to show the H-C-H bond angle is 109.5°

Graphene	 <i>Single layer of graphite one atom thick</i>	Excellent conductor.	Contains delocalised electrons.
		Very strong.	Contains strong covalent bonds.


Graphene and fullerenes	 Buckminsterfullerene, C ₆₀ First fullerene to be discovered.	Hexagonal rings of carbon atoms with hollow shapes. Can also have rings of five (pentagonal) or seven (heptagonal) carbon atoms.
-------------------------	---	--

Use of nanoparticles	<i>Healthcare, cosmetics, sun cream, catalysts, deodorants, electronics.</i>	Nanoparticles may be toxic to people. They may be able to enter the brain from the bloodstream and cause harm.
----------------------	--	--

Fullerenes		Very conductive.	Used in electronics industry.
		<i>Very thin and long cylindrical fullerenes</i>	Reinforcing composite materials.
		High tensile strength.	Catalysts and lubricants.

Carbon nanotubes		Very thin and long cylindrical fullerenes	Used in electronics industry.
		High tensile strength.	Reinforcing composite materials.
		Large surface area to volume ratio.	Catalysts and lubricants.

Each carbon atom is bonded to three others forming layers of hexagonal rings with no covalent bonds between the layers



Graphite	
Slippery.	Layers can slide over each other.
Very high melting point.	Strong covalent bonds.
Does conduct electricity.	Delocalised electrons between layers.

Oxidation Is Loss (of electrons) **Reduction Is Gain** (of electrons)

HT ONLY: Reactions between metals and acids are redox reactions as the metal donates electrons to the hydrogen ions. This displaces hydrogen as a gas while the metal ions are left in the solution.

Ionic half equations (HT only)

For displacement reactions

Ionic half equations show what happens to each of the reactants during reactions

For example:
The ionic equation for the reaction between iron and copper (II) ions is:
 $Fe + Cu^{2+} \rightarrow Fe^{2+} + Cu$

The half-equation for iron (II) is:
 $Fe \rightarrow Fe^{2+} + 2e^{-}$

The half-equation for copper (II) ions is:
 $Cu^{2+} + 2e^{-} \rightarrow Cu$

Reactions with acids

metal + acid → salt + hydrogen

magnesium + hydrochloric acid → magnesium chloride + hydrogen

zinc + sulfuric acid → zinc sulfate + hydrogen

Acids react with some metals to produce salts and hydrogen.

Extraction using carbon

Metals less reactive than carbon can be extracted from their oxides by reduction.

For example:
zinc oxide + carbon → zinc + carbon dioxide

Reactions of acids and metals

Reactions of acids

Extraction of metals and reduction

Unreactive metals, such as gold, are found in the Earth as the metal itself. They can be mined from the ground.

AQA Chemical Changes 1

Reactivity of metals

	Reactions with water	Reactions with acid
Group 1 metals	<i>Reactions get more vigorous as you go down the group</i>	<i>Reactions get more vigorous as you go down the group</i>
Group 2 metals	<i>Do not react with water</i>	<i>Observable reactions include fizzing and temperature increases</i>
Zinc, iron and copper	<i>Do not react with water</i>	<i>Zinc and iron react slowly with acid. Copper does not react with acid.</i>

The reactivity series

Metals form positive ions when they react	<i>The reactivity of a metal is related to its tendency to form positive ions</i>	The reactivity series arranges metals in order of their reactivity (their tendency to form positive ions).
Carbon and hydrogen	<i>Carbon and hydrogen are non-metals but are included in the reactivity series</i>	These two non-metals are included in the reactivity series as they can be used to extract some metals from their ores, depending on their reactivity.
Displacement	<i>A more reactive metal can displace a less reactive metal from a compound.</i>	Silver nitrate + Sodium chloride → Sodium nitrate + Silver chloride

- potassium **most reactive** K
sodium Na
calcium Ca
magnesium Mg
aluminium Al
carbon C
zinc Zn
iron Fe
tin Sn
lead Pb
hydrogen H
copper Cu
silver Ag
gold Au
platinum **least reactive** Pt
-

Oxidation and reduction in terms of electrons (HT ONLY)

Neutralisation of acids and salt production

Acid name	Salt name
<i>Hydrochloric acid</i>	Chloride
<i>Sulfuric acid</i>	Sulfate
<i>Nitric acid</i>	Nitrate

sodium hydroxide + hydrochloric acid → sodium chloride + water

calcium carbonate + sulfuric acid → calcium sulfate, + carbon dioxide + water

Neutralisation

Acids can be neutralised by alkalis and bases

An **alkali** is a soluble base e.g. metal hydroxide.
A **base** is a substance that neutralises an acid e.g. a soluble metal hydroxide or a metal oxide.

Metals and oxygen	<i>Metals react with oxygen to form metal oxides</i>	magnesium + oxygen → magnesium oxide $2Mg + O_2 \rightarrow 2MgO$
Reduction	<i>This is when oxygen is removed from a compound during a reaction</i>	e.g. metal oxides reacting with hydrogen, extracting low reactivity metals
Oxidation	<i>This is when oxygen is gained by a compound during a reaction</i>	e.g. metals reacting with oxygen, rusting of iron

Metal oxides

The ions discharged when an aqueous solution is electrolysed using inert electrodes depend on the relative reactivity of the elements involved.

Electrolysis of aqueous solutions

At the negative electrode	Metal will be produced on the electrode if it is less reactive than hydrogen. Hydrogen will be produced if the metal is more reactive than hydrogen.
At the positive electrode	Oxygen is formed at positive electrode. If you have a halide ion (Cl ⁻ , I ⁻ , Br ⁻) then you will get chlorine, bromine or iodine formed at that electrode.

Process of electrolysis	<i>Splitting up using electricity</i>	When an ionic compound is melted or dissolved in water, the ions are free to move. These are then able to conduct electricity and are called electrolytes. Passing an electric current through electrolytes causes the ions to move to the electrodes.
Electrode	Anode Cathode	The positive electrode is called the anode. The negative electrode is called the cathode.
Where do the ions go?	Cations Anions	Cations are positive ions and they move to the negative cathode. Anions are negative ions and they move to the positive anode.

Extracting metals using electrolysis	<i>Metals can be extracted from molten compounds using electrolysis.</i>
	<i>This process is used when the metal is too reactive to be extracted by reduction with carbon.</i>
	<i>The process is expensive due to large amounts of energy needed to produce the electrical current. Example: aluminium is extracted in this way.</i>
	<i>Aluminium extraction uses CRYOLITE to lower the melting point of aluminium oxide. Electrodes have to be replaced regularly as they are made of GRAPHITE (CARBON) which react with the oxygen produced.</i>

Strong acids	<i>Completely ionised in aqueous solutions e.g. hydrochloric, nitric and sulfuric acids.</i>
Weak acids	<i>Only partially ionised in aqueous solutions e.g. ethanoic acid, citric acid.</i>
Hydrogen ion concentration	<i>As the pH decreases by one unit (becoming a stronger acid), the hydrogen ion concentration increases by a factor of 10.</i>

Electrolysis

AQA Chemical Changes 2

Higher tier: Half equations, for example:
 At the cathode: $Pb^{2+} + 2e^{-} \rightarrow Pb$
 At the anode: $2Br^{-} \rightarrow Br_2 + 2e^{-}$


Soluble salts	<i>Soluble salts can be made from reacting acids with solid insoluble substances (e.g. metals, metal oxides, hydroxides and carbonates).</i>
Production of soluble salts	<i>Add the solid to the acid until no more dissolves. Filter off excess solid and then crystallise to produce solid salts.</i>

Reactions of acids

Titrations (Chemistry only)

Titration is used to work out the precise volumes of acid and alkali solutions that react with each other.

- Use the pipette to add 25 cm³ of alkali to a conical flask and add a few drops of indicator. (a **pipette** is used for **fixed volumes** only)
- Fill the burette with acid and note the starting volume. Slowly add the acid from the burette to the alkali in the conical flask, swirling to mix. (a **burette** is used for **variable volumes**)
- Stop adding the acid when the end-point is reached (the appropriate colour change in the indicator happens). Note the final volume reading. Repeat steps 1 to 3 until you get consistent readings.



You can use universal indicator or a pH probe to measure the acidity or alkalinity of a solution against the pH scale.

The pH scale and neutralisation

In neutralisation reactions, hydrogen ions react with hydroxide ions to produce water:
 $H^{+}_{(aq)} + OH^{-}_{(aq)} \rightarrow H_2O_{(l)}$

Acids	<i>Acids contain hydrogen ions (H⁺) in aqueous solutions.</i>
Alkalis	<i>Aqueous solutions of alkalis contain hydroxide ions (OH⁻).</i>

Calculating the chemical quantities in titrations involving concentrations in mol/dm³ and in g/dm³ (HT ONLY):

$2NaOH(aq) + H_2SO_4(aq) \rightarrow Na_2SO_4(aq) + 2H_2O(l)$

It takes 12.20cm³ of sulfuric acid to neutralise 24.00cm³ of sodium hydroxide solution, which has a concentration of 0.50mol/dm³.

Calculate the concentration of the sulfuric acid in g/dm³
 $0.5 \text{ mol/dm}^3 \times (24/1000) \text{ dm}^3 = 0.012 \text{ mol of NaOH}$

The equation shows that 2 mol of NaOH reacts with 1 mol of H₂SO₄, so the number of moles in 12.20cm³ of sulfuric acid is **(0.012/2) = 0.006 mol of sulfuric acid**

Calculate the concentration of sulfuric acid in mol/dm³
 $0.006 \text{ mol} \times (1000/12.2) \text{ dm}^3 = 0.49 \text{ mol/dm}^3$

Calculate the concentration of sulfuric acid in g/dm³
 $H_2SO_4 = (2 \times 1) + 32 + (4 \times 16) = 98 \text{g}$
 $0.49 \times 98 = 48.2 \text{g/dm}^3$

AQA GCSE QUANTITATIVE CHEMISTRY

M_r	<i>The sum of the relative atomic masses of the atoms in the numbers shown in the formula. The M_r is the mass of 1 mole of the formula in grams.</i>	The sum of the M_r of the reactants in the quantities shown equals the sum of the M_r of the products in the quantities shown.	$2Mg + O_2 \rightarrow 2MgO$ $48g + 32g = 80g$ $80g = 80g$
-------	--	--	--

The reactant that is completely used up	<i>Limits the amount of product that is made</i>	Less moles of product are made.
---	--	---------------------------------

Balanced symbol equations	<i>Represent chemical reactions and have the same number of atoms of each element on both sides of the equation</i>	<p style="text-align: center;">$H_2 + Cl_2 \rightarrow 2HCl$</p> <p style="text-align: center;">Subscript Normal script</p> <p>Subscript (small number at the bottom, after the element) numbers show the number of atoms of the element to its left.</p> <p>Normal script (in front of the formula) numbers show the number of molecules.</p>
---------------------------	---	--

Conservation of mass and balanced symbol equations

Relative formula mass (M_r)

Limiting reactants (HT only)

Chemical measurements

Whenever a measurement is taken, there is always some uncertainty about the result obtained	<i>Can determine whether the mean value falls within the range of uncertainty of the result</i>	<ol style="list-style-type: none"> 1. Calculate the mean 2. Calculate the range of the results 3. Estimate of uncertainty in mean would be half the range
---	---	--

Conservation of mass	<i>No atoms are lost or made during a chemical reaction</i>	Mass of the products equals the mass of the reactants.
----------------------	---	--

Concentration of solutions		
Measured in mass per given volume of solution (g/dm^3)	<i>Conc. = $\frac{mass (g)}{volume (dm^3)}$</i>	HT only Greater mass = higher concentration. Greater volume = lower concentration.

Mass changes when a reactant or product is a gas		
Mass appears to increase during a reaction	<i>One of the reactants is a gas</i>	Magnesium + oxygen \rightarrow magnesium oxide
Mass appears to decrease during a reaction	<i>One of the products is a gas and has escaped</i>	Calcium carbonate \rightarrow carbon dioxide + calcium oxide

Moles (HT only)

Amounts of substances in equations (HT only)

Using moles to balance equations (HT only)

The balancing numbers in a symbol equation can be calculated from the masses of reactants and products	<i>Convert the masses in grams to amounts in moles and convert the number of moles to simple whole number ratios.</i>
--	---

Chemical amounts are measured in moles (mol)	<i>Mass of one mole of a substance in grams = relative formula mass</i>	One mole of $H_2O = 18g (1 + 1 + 16)$ One mole of $Mg = 24g$
--	---	---

Avogadro constant	6.02×10^{23} per mole One mole of H_2O will contain 6.02×10^{23} molecules One mole of $NaCl$ will contain 6.02×10^{23} Na^+ ions
-------------------	--

<i>Number of moles = $\frac{mass (g)}{A_r}$ or $\frac{mass (g)}{M_r}$</i>	How many moles of sulfuric acid molecules are there in 4.7g of sulfuric acid (H_2SO_4)? Give your answer to 1 significant figure. $\frac{4.7}{98} = 0.05 \text{ mol}$ (M_r of H_2SO_4)
---	---

Chemical equations show the number of moles reacting and the number of moles made

$Mg + 2HCl \rightarrow MgCl_2 + H_2$
One mole of magnesium reacts with two moles of hydrochloric acid to make one mole of magnesium chloride and one mole of hydrogen

If you have a 60g of Mg, what mass of HCl do you need to convert it to $MgCl_2$? $A_r : Mg = 24$ so mass of 1 mole of $Mg = 24g$ $M_r : HCl (1 + 35.5)$ so mass of 1 mole of $HCl = 36.5g$ So 60g of Mg is $60/24 = 2.5$ moles Balanced symbol equation tells us that for every one mole of Mg , you need two moles of HCl to react with it. So you need $2.5 \times 2 = 5$ moles of HCl You will need $5 \times 36.5g$ of $HCl = 182.5g$
--


A measure of the amount of starting materials that end up as useful products

Atom economy = $\frac{\text{Relative formula mass of desired product from equation}}{100} \times 100$

Sum of relative formula mass of all reactants from equation

High atom economy is important for sustainable development and economic reasons

Calculate the atom economy for making hydrogen by reacting zinc with hydrochloric acid:



$$M_r \text{ of } \text{H}_2 = 1 + 1 = 2$$

$$M_r \text{ of } \text{Zn} + 2\text{HCl} = 65 + 1 + 1 + 35.5 + 35.5 = 138$$

$$\text{Atom economy} = \frac{2}{138} \times 100 = \frac{2}{138} \times 100 = 1.45\%$$

This method is unlikely to be chosen as it has a low atom economy.

Atom economy

Concentration of a solution is the amount of solute per volume of solution

$$\text{Concentration} = \frac{\text{moles (mol)}}{\text{volume (dm}^3\text{)}} \text{ (mol/dm}^3\text{)}$$

$$\text{Concentration} = \frac{\text{mass (g)}}{\text{volume (dm}^3\text{)}} \text{ (g/dm}^3\text{)}$$

$$\text{g/dm}^3 \xrightleftharpoons[\times M_r]{\div M_r} \text{mol/dm}^3$$

What is the concentration of a solution that has 35.0g of solute in 0.5dm³ of solution?


$$35/0.5 = 70 \text{ g/dm}^3$$

Using concentrations of solutions in mol/dm³ (HT only, chemistry only)

AQA QUANTITATIVE CHEMISTRY 2

Titration

If the volumes of two solutions that react completely are known and the concentrations of one solution is known, the concentration of the other solution can be calculated.



It takes 12.20cm³ of sulfuric acid to neutralise 24.00cm³ of sodium hydroxide solution, which has a concentration of 0.50mol/dm³.

Calculate the concentration of the sulfuric acid in mol/dm³:

$$0.5 \text{ mol/dm}^3 \times (24/1000) \text{ dm}^3 = 0.012 \text{ mol of NaOH}$$

The equation shows that 2 mol of NaOH reacts with 1 mol of H₂SO₄, so the number of moles in 12.20cm³ of sulfuric acid is (0.012/2) = 0.006 mol of sulfuric acid

$$\text{Calculate the concentration of sulfuric acid in mol/dm}^3: 0.006 \text{ mol} \times (1000/12.2) \text{ dm}^3 = 0.49 \text{ mol/dm}^3$$

HT only:

200g of calcium carbonate is heated. It decomposes to make calcium oxide and carbon dioxide. Calculate the theoretical mass of calcium oxide made.



$$M_r \text{ of } \text{CaCO}_3 = 40 + 12 + (16 \times 3) = 100$$

$$M_r \text{ of } \text{CaO} = 40 + 16 = 56$$

$$100\text{g of } \text{CaCO}_3 \text{ would make } 56 \text{ g of } \text{CaO}$$

$$\text{So } 200\text{g would make } 112\text{g}$$

Percentage yield

Use of amount of substance in relation to volumes of gases (HT only, chemistry only)

Calculate the concentration of sulfuric acid in g/dm³:

$$\text{H}_2\text{SO}_4 = (2 \times 1) + 32 + (4 \times 16) = 98\text{g} \\ 0.49 \times 98\text{g} = 48.2\text{g/dm}^3$$

Yield is the amount of product obtained

It is not always possible to obtain the calculated amount of a product

The reaction may not go to completion because it is reversible.

Some of the product may be lost when it is separated from the reaction mixture.

Some of the reactants may react in ways different to the expected reaction.

Equal amounts of moles or gases occupy the same volume under the same conditions of temperature and pressure

The volume of one mole of any gas at room temperature and pressure (20°C and 1 atmospheric pressure) is **24 dm³**

No. of moles of gas = $\frac{\text{vol of gas (dm}^3\text{)}}{24\text{dm}^3}$

$$24\text{dm}^3$$

Percentage yield is comparing the amount of product obtained as a percentage of the maximum theoretical amount

$$\% \text{ Yield} = \frac{\text{Mass of product made} \times 100}{\text{Max. theoretical mass}}$$

A piece of sodium metal is heated in chlorine gas. A maximum theoretical mass of 10g for sodium chloride was calculated, but the actual yield was only 8g.

Calculate the percentage yield.

$$\text{Percentage yield} = 8/10 \times 100 = 80\%$$

What is the volume of 11.6 g of butane (C₄H₁₀) gas at RTP?

$$M_r : (4 \times 12) + (10 \times 1) = 58$$

$$11.6/58 = 0.20 \text{ mol}$$

$$\text{Volume} = 0.20 \times 24 = 4.8 \text{ dm}^3$$

6g of a hydrocarbon gas had a volume of 4.8 dm³. Calculate its molecular mass.

$$1 \text{ mole} = 24 \text{ dm}^3, \text{ so } 4.8/24 = 0.2 \text{ mol}$$

$$M_r = 6 / 0.2 = 30$$


$$\text{If } 6\text{g} = 0.2 \text{ mol}, 1 \text{ mol equals } 30 \text{ g}$$

Rate of chemical reaction	<i>This can be calculated by measuring the quantity of reactant used or product formed in a given time.</i>	Rate = $\frac{\text{quantity of reactant used}}{\text{time taken}}$
		Rate = $\frac{\text{quantity of product formed}}{\text{time taken}}$

Factors affecting the rate of reaction	
Temperature	<i>The higher the temperature, the quicker the rate of reaction.</i>
Concentration	<i>The higher the concentration, the quicker the rate of reaction.</i>
Surface area	<i>The larger the surface area of a reactant solid, the quicker the rate of reaction.</i>
Pressure (of gases)	<i>When gases react, the higher the pressure upon them, the quicker the rate of reaction.</i>

Quantity	Unit
Mass	Grams (g)
Volume	cm ³
Rate of reaction	Grams per cm ³ (g/cm ³) HT: moles per second (mol/s)


Calculating rates of reactions



Rate of reaction

Factors affecting rates


Collision theory and activation energy



Collision theory	<i>Chemical reactions can only occur when reacting particles collide with each other with sufficient energy.</i>	Increasing the temperature increases the frequency of collisions and makes the collisions more energetic, therefore increasing the rate of reaction.
Activation energy	<i>This is the minimum amount of energy colliding particles in a reaction need in order to react.</i>	Increasing the concentration, pressure (gases) and surface area (solids) of reactions increases the frequency of collisions, therefore increasing the rate of reaction.

AQA GCSE The rate and extent of chemical change

Reversible reactions and dynamic equilibrium



Catalyst	A catalyst changes the rate of a chemical reaction but is not used in the reaction.
Enzymes	These are biological catalysts.
How do they work?	Catalysts provide a different reaction pathway where reactants do not require as much energy to react when they collide

If a catalyst is used in a reaction, it is not shown in the word equation.


Catalysts

Changing conditions and equilibrium (HT)

The relative amounts of reactants and products at equilibrium depend on the conditions of the reaction.

Le Chatelier's Principles	States that when a system experiences a disturbance (change in condition), it will respond to restore a new equilibrium state.
Changing concentration	If the concentration of a reactant is increased, more products will be formed. If the concentration of a product is decreased, more reactants will react.
Changing temperature	If the temperature of a system at equilibrium is increased: - Exothermic reaction = products decrease - Endothermic reaction = products increase
Changing pressure (gaseous reactions)	For a gaseous system at equilibrium: - Pressure increase = equilibrium position shifts to side of equation with smaller number of molecules. - Pressure decrease = equilibrium position shifts to side of equation with larger number of molecules.

Reversible reactions
In some chemical reactions, the products can react again to re-form the reactants



The direction
The direction of reversible reactions can be changed by changing conditions:


$$A + B \xrightleftharpoons[\text{cool}]{\text{heat}} C + D$$

Energy changes and reversible reactions

If one direction of a reversible reaction is exothermic, the opposite direction is endothermic. The same amount of energy is transferred in each case.

Equilibrium

Equilibrium in reversible reactions
When a reversible reaction occurs in apparatus which prevents the escape of reactants and products, equilibrium is reached when the forward and reverse reactions occur exactly at the same rate.



Endothermic	<i>Energy is taken in from the surroundings so the temperature of the surroundings decreases</i>	<ul style="list-style-type: none"> Thermal decomposition Sports injury packs
Exothermic	<i>Energy is transferred to the surroundings so the temperature of the surroundings increases</i>	<ul style="list-style-type: none"> Combustion Hand warmers Neutralisation

Ionic half equations	Negative electrode: $2\text{H}_2(\text{g}) + 4\text{OH}^-(\text{aq}) \rightarrow 4\text{H}_2\text{O}(\text{l}) + 4\text{e}^-$	Positive electrode: $\text{O}_2(\text{g}) + 2\text{H}_2\text{O}(\text{l}) + 4\text{e}^- \rightarrow 4\text{OH}^-(\text{aq})$
Hydrogen fuel cells	Word equation: <i>hydrogen + oxygen → water</i>	Symbol equation: $2\text{H}_2 + \text{O}_2 \rightarrow 2\text{H}_2\text{O}$
	Advantages: <ul style="list-style-type: none"> No pollutants produced Can be a range of sizes 	Disadvantages: <ul style="list-style-type: none"> Hydrogen is highly flammable Hydrogen is difficult to store

Reaction profiles	<i>Show the overall energy change of a reaction</i>
-------------------	---

Breaking bonds in reactants	<i>Endothermic process</i>
Making bonds in products	<i>Exothermic process</i>

Overall energy change of a reaction	<i>Exothermic</i>	Energy released making new bonds is greater than the energy taken in breaking existing bonds.
	<i>Endothermic</i>	Energy needed to break existing bonds is greater than the energy released making new bonds.

Types of reaction

Fuel cells (Chemistry only)

The energy change of reactions (HT only)

AQA GCSE Energy changes

Reaction profiles

Activation energy	<i>Chemical reactions only happen when particles collide with sufficient energy</i>	The minimum amount of energy that colliding particles must have in order to react is called the activation energy.
-------------------	---	--

Cells and batteries (Chemistry only)

Simple cell	<i>Make a simple cell by connecting two different metals in contact with an electrolyte</i>	Increase the voltage by increasing the reactivity difference between the two metals.
Batteries	<i>Consist of two or more cells connected together in series to provide a greater voltage.</i>	
Non-rechargeable cells	<i>Stop when one of the reactants has been used up</i>	Alkaline batteries
Rechargeable cells	<i>Can be recharged because the chemical reactions are reversed when an external electrical current is supplied</i>	Rechargeable batteries

Endothermic		Products are at a higher energy level than the reactants. As the reactants form products, energy is transferred from the surroundings to the reaction mixture. The temperature of the surroundings decreases because energy is taken in during the reaction.
Exothermic		Products are at a lower energy level than the reactants. When the reactants form products, energy is transferred to the surroundings. The temperature of the surroundings increases because energy is released during the reaction.

Bond energy calculation	Calculate the overall energy change for the forward reaction $\text{N}_2 + 3\text{H}_2 \rightleftharpoons 2\text{NH}_3$
	Bond energies (in kJ/mol): H-H 436, H-N 391, N≡N 945
	Bond breaking: $945 + (3 \times 436) = 945 + 1308 = 2253 \text{ kJ/mol}$
	Bond making: $6 \times 391 = 2346 \text{ kJ/mol}$
	Overall energy change = $2253 - 2346 = -93 \text{ kJ/mol}$
	Therefore reaction is exothermic overall.